arXiv:0805.3679v2 [hep-th] 19 Apr 2009

Nonlinear Schrödinger equations and N=2 superconformal algebra

H. T.Özer *

Physics Department,
Faculty of Science and Letters,
Istanbul Technical University,
34469, Maslak, Istanbul, Turkey

April 19, 2009

We obtain new coupled super Nonlinear Schrödinger equations by using AKNS scheme and soliton connection taking values in N=2 superconformal algebra .

1 Introduction

Coupled Nonlinear Schrödinger (NLS) equations can be obtained using Ablowitz, Kaub, Newell , Segur (AKNS) scheme[1].Extensions of coupled NLS equations have been obtained using a simple Lie algebra[2], a Kac-Moody algebra [3],a Lie superalgebra[3,4],a Virasoro algebra[5] and a N=1 superconformal algebra [6]in the literature.

The N=2 superconformal algebras were discovered in the seventies independently by Ademollo et al.[7] and by Kac[8].The first authors derived the algebras for physical purposes, in order to define supersymmetric strings, whereas Kac derived them for mathematical purposes along with his classification of Lie superalgebras.

The N=2 superconformal algebras provide the symmetries underlying the N=2 strings[9,10,11,12,13]. These seem to be related to M-theory since many of the basic objects of M-theory are realized in the heterotic (2,1) N=2 strings[14]. In addition, the topological version of the N=2 superconformal algebra is realized in the world-sheet of the bosonic string[15],as well as in the world-sheet of the superstrings[16].

In literature super-extensions of Nonlinear Schroedinger equations involve finite number of bosonic and finite number of fermionic fields [3,4,17,18,19].The super-extensions of Nonlinear Schroedinger equations with infinite number of bosonic fields and infinite number of fermionic fields has been studied in ref.6 and in this paper.Mathematically this paper contains extension of the work done in ref.6.

In this paper we will obtain super - extensions of coupled NLS equations using N=2 superconformal algebra with Neveu-Schwarz and Ramond types . In sec.2 we will discuss the sl(2,1)superalgebra valued soliton connection and we will obtain coupled super NLS equations. Sec.3 and sec.4 concern the soliton connection for the N=2 superconformal algebra with Neveu-Schwarz type and Ramond type, respectively and we will obtain in these sections two different types of super-extensions of coupled NLS equations.

---

*Corresponding author. **E-mail : ozert @ itu.edu.tr**

2 AKNS Scheme with sl(2,1) Superalgebra

In AKNS scheme in 1+1 dimension the connection is defined as

$$\Omega = \Omega_b + \Omega_f \tag{1}$$

where

$$\Omega_b = \Big(i\lambda J_1 + i\lambda H_1 + Q^{+1}E_{+1} + Q^{-1}E_{-1}\Big)dx + \Big(- A_1 J_1 - A_2 H_1 + B^{+1}E_{+1} + B^{-1}E_{-1} + \Big)dt \tag{2}$$

$$\Omega_f = \Big(P_1^{+\frac{1}{2}}F_{+\frac{1}{2}} + P_1^{-\frac{1}{2}}F_{-\frac{1}{2}} + P_2^{+\frac{1}{2}}\bar{F}_{+\frac{1}{2}} + P_2^{-\frac{1}{2}}\bar{F}_{-\frac{1}{2}}\Big)dx + \Big(C_1^{+\frac{1}{2}}F_{+\frac{1}{2}} + C_1^{-\frac{1}{2}}F_{-\frac{1}{2}} + C_2^{+\frac{1}{2}}\bar{F}_{+\frac{1}{2}} + C_2^{-\frac{1}{2}}\bar{F}_{-\frac{1}{2}}\Big)dt \tag{3}$$

where J_1,H_1, $E_{\pm1}$are bosonic generators and $F_{\pm\frac{1}{2}}\bar{F}_{\pm\frac{1}{2}}$ are fermionic generators of $sl(1,2) \cong sl2,1)$ superalgebra which is the (N=2)extended supersymmetric version of sl(2) algebra. These generators have matrix representations as

$$\begin{gathered}
H_1 = \begin{pmatrix} \frac{1}{2} & 0 & 0 \\ 0 & -\frac{1}{2} & 0 \\ 0 & 0 & 0 \end{pmatrix}; J_1 = \begin{pmatrix} \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \\ 0 & 0 & 0 \end{pmatrix}; E_{+1} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}; E_{-1} = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\
F_{+\frac{1}{2}} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}; F_{-\frac{1}{2}} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \bar{F}_{+\frac{1}{2}} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}; \bar{F}_{-\frac{1}{2}} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}
\end{gathered} \tag{4}$$

Also, these generators satisfy the following commutation and anticommutation relations

$$\begin{array}{ccc}
[H_1, E_{\pm1}] = \pm E_{\pm1}; & [H_1, F_{\pm1}] = \pm\frac{1}{2}F_{\pm1}; & \Big[H_1, \bar{F}_{\pm1}\Big] = \pm\frac{1}{2}\bar{F}_{\pm1} \\
[J_1, H_{\pm1}] = [J_1, E_{\pm1}] = 0; & [J_1, F_{\pm1}] = \frac{1}{2}F_{\pm1}; & \Big[J_1, \bar{F}_{\pm1}\Big] = -\frac{1}{2}\bar{F}_{\pm1} \\
\Big[E_{\pm1}, F_{\mp\frac{1}{2}}\Big] = \Big[E_{\pm1}, \bar{F}_{\mp\frac{1}{2}}\Big] = 0; & \Big[E_{\pm1}, F_{\mp\frac{1}{2}}\Big] = -F_{\pm\frac{1}{2}}; & \Big[E_{\pm1}, \bar{F}_{\mp\frac{1}{2}}\Big] = \bar{F}_{\pm\frac{1}{2}} \\
\Big\{F_{\pm1}, F_{\pm\frac{1}{2}}\Big\} = \Big\{\bar{F}_{\pm1}, \bar{F}_{\pm\frac{1}{2}}\Big\} = 0; & \Big\{F_{\pm1}, F_{\mp\frac{1}{2}}\Big\} = \Big\{\bar{F}_{\pm1}, \bar{F}_{\mp\frac{1}{2}}\Big\} = 0; & \Big\{F_{\pm1}, \bar{F}_{\mp\frac{1}{2}}\Big\} = E_{\pm1} \\
[E_{+1}, E_{-1}] = H_1; & \Big\{F_{\pm\frac{1}{2}}, \bar{F}_{\pm\frac{1}{2}}\Big\} = J_1 \mp H_1 &
\end{array} \tag{5}$$

In Eq.(1) λ is the spectral parameter, $Q^{\pm1}$,$P_1^{\pm\frac{1}{2}}$ and $P_2^{\pm\frac{1}{2}}$ are fields depending on space and time, namely x and t, and functions A_1,A_2,$B^{\pm1}$,$C_1^{\pm\frac{1}{2}}$ and $C_2^{\pm\frac{1}{2}}$ are x,t and λ dependent. The integrability condition is given by

$$d\,\Omega \;+\; \Omega \;\wedge\; \Omega = 0 \tag{6}$$

By using Eqs.(1) and (6) one can obtain following equations:

$$A_{1x} = P_1^{-\frac{1}{2}}C_2^{+\frac{1}{2}} + P_1^{+\frac{1}{2}}C_2^{-\frac{1}{2}} + P_2^{-\frac{1}{2}}C_1^{+\frac{1}{2}} + P_2^{+\frac{1}{2}}C_1^{-\frac{1}{2}} \tag{7}$$

$$A_{2x} = -2Q^{-1}B^{+1} + 2Q^{+1}B^{-1} + P_1^{-\frac{1}{2}}C_2^{+\frac{1}{2}} - P_1^{+\frac{1}{2}}C_2^{-\frac{1}{2}} - P_2^{-\frac{1}{2}}C_1^{+\frac{1}{2}} + P_2^{+\frac{1}{2}}C_1^{-\frac{1}{2}} \tag{8}$$

$$Q^{+1}{}_t = \quad B^{+1}{}_x + i\lambda B^{+1} + Q^{+1}A_2 + P_1^{+\frac{1}{2}}C_2^{+\frac{1}{2}} + P_2^{+\frac{1}{2}}C_1^{+\frac{1}{2}} \tag{9}$$

$$Q^{-1}{}_t = \quad B^{-1}{}_x - i\lambda B^{-1} - Q^{-1}A_2 + P_1^{-\frac{1}{2}}C_2^{-\frac{1}{2}} + P_2^{-\frac{1}{2}}C_1^{-\frac{1}{2}} \tag{10}$$

$$P_1^{+\frac{1}{2}}{}_t = \quad C_1^{+\frac{1}{2}}{}_x + i\lambda C_1^{+\frac{1}{2}} + P_1^{-\frac{1}{2}}B^{+1} + \tfrac{1}{2}P_1^{+\frac{1}{2}}A_1 + \tfrac{1}{2}P_1^{+\frac{1}{2}}A_2 - Q^{+1}C_1^{-\frac{1}{2}} \tag{11}$$

$$P_1^{-\frac{1}{2}}{}_t = \quad C_1^{-\frac{1}{2}}{}_x + P_1^{+\frac{1}{2}}B^{-1} + \tfrac{1}{2}P_1^{-\frac{1}{2}}A_1 - \tfrac{1}{2}P_1^{-\frac{1}{2}}A_2 - Q^{-1}C_1^{+\frac{1}{2}} \tag{12}$$

$$P_2^{+\frac{1}{2}}{}_t = \quad C_2^{+\frac{1}{2}}{}_x - P_2^{-\frac{1}{2}}B^{+1} - \tfrac{1}{2}P_2^{+\frac{1}{2}}A_1 + \tfrac{1}{2}P_2^{+\frac{1}{2}}A_2 + Q^{+1}C_2^{-\frac{1}{2}} \tag{13}$$

$$P_2^{-\frac{1}{2}}{}_t = \quad C_2^{-\frac{1}{2}}{}_x - i\lambda C_2^{-\frac{1}{2}} - P_2^{+\frac{1}{2}}B^{-1} - \tfrac{1}{2}P_2^{-\frac{1}{2}}A_1 - \tfrac{1}{2}P_2^{-\frac{1}{2}}A_2 + Q^{-1}C_2^{+\frac{1}{2}} \tag{14}$$

In AKNS scheme we expand A_1,A_2,$B^{\pm 1}$ $C_1^{\pm\frac{1}{2}}$ and $C_2^{\pm\frac{1}{2}}$ in terms of positive powers of λ as

$$A_1 = \sum_{n=0}^{2} \lambda^n a_{1n}; \quad A_2 = \sum_{n=0}^{2} \lambda^n a_{2n}; \quad B^{\pm 1} = \sum_{n=0}^{2} \lambda^n b_n^{\pm 1}; \quad C_1^{\pm\frac{1}{2}} = \sum_{n=0}^{2} \lambda^n c_{1n}^{\pm\frac{1}{2}}; \quad C_2^{\pm\frac{1}{2}} = \sum_{n=0}^{2} \lambda^n c_{2n}^{\pm\frac{1}{2}} \tag{15}$$

Inserting Eq.(15) into Eqs.(7-14)gives 24 relations in terms of a_{1n},a_{2n},$b_n^{\pm 1}$,$c_{1n}^{\pm\frac{1}{2}}$ and $c_{2n}^{\pm\frac{1}{2}}$. By solving these relations we get

$$\begin{gathered}
a_{10} = -iP_1^{-\frac{1}{2}} P_2^{+\frac{1}{2}} - iP_1^{+\frac{1}{2}} P_2^{-\frac{1}{2}}; \quad a_{11} = 0; \quad a_{12} = -i; \\
a_{20} = -2iQ^{+1}Q^{-1} - iP_1^{-\frac{1}{2}} P_2^{+\frac{1}{2}} + iP_1^{+\frac{1}{2}} P_2^{-\frac{1}{2}}; \quad a_{21} = 0; \quad a_{22} = -i; \\
b_0^{\pm 1} = \pm i{Q^{\pm 1}}_x + iP_2^{\pm\frac{1}{2}} P_1^{\pm\frac{1}{2}}; \; b_1^{\pm 1} = Q^{\pm 1}; \; b_2^{\pm 1} = 0 \\
c_{10}^{\pm\frac{1}{2}} = iP_1^{\mp\frac{1}{2}} Q^{\pm 1} + i{P_1^{\pm}}_x; \; c_{20}^{\pm\frac{1}{2}} = iP_2^{\mp\frac{1}{2}} Q^{\pm 1} - i{P_2^{\pm}}_x \\
c_{11}^{+\frac{1}{2}} = P^{+\frac{1}{2}}; \; c_{21}^{-\frac{1}{2}} = P^{-\frac{1}{2}}; \; c_{11}^{-\frac{1}{2}} = c_{21}^{+\frac{1}{2}} = 0; \; c_{12}^{\pm\frac{1}{2}} = c_{22}^{\pm\frac{1}{2}} = 0
\end{gathered} \tag{16}$$

By using the relations given by Eq.(16) from Eqs.(9-14) we obtain the coupled super NLS equations as

$$\begin{aligned}
-i{Q^{+1}}_t &= {Q^{+1}}_{xx} - 2(Q^{+1})^2 Q^{-1} + 2Q^{+1} P_2^{+\frac{1}{2}} P_1^{-\frac{1}{2}} - 2Q^{+1} P_2^{-\frac{1}{2}} P_1^{+\frac{1}{2}} + 2(P_2^{+\frac{1}{2}} P_1^{+\frac{1}{2}})_x \\
i{Q^{-1}}_t &= {Q^{+1}}_{xx} - 2(Q^{-1})^2 Q^{+1} - 2Q^{-1} P_2^{-\frac{1}{2}} P_1^{+\frac{1}{2}} + 2Q^{-1} P_2^{+\frac{1}{2}} P_1^{-\frac{1}{2}} - 2(P_2^{-\frac{1}{2}} P_1^{-\frac{1}{2}})_x \\
-i{P_1^{+\frac{1}{2}}}_t &= {P_1^{+\frac{1}{2}}}_{xx} - 2P_1^{+\frac{1}{2}} Q^{+1} Q^{-1} + 2P_1^{-\frac{1}{2}} {Q^{+1}}_x - 2P_2^{+\frac{1}{2}} P_1^{+\frac{1}{2}} P_1^{-\frac{1}{2}} \\
-i{P_1^{-\frac{1}{2}}}_t &= {P_1^{-\frac{1}{2}}}_{xx} - 2P_1^{-\frac{1}{2}} Q^{+1} Q^{-1} - 2P_2^{-\frac{1}{2}} P_1^{-\frac{1}{2}} P_1^{+\frac{1}{2}} \\
i{P_2^{+\frac{1}{2}}}_t &= {P_2^{+\frac{1}{2}}}_{xx} + 2P_2^{+\frac{1}{2}} Q^{+1} Q^{-1} + 2P_2^{+\frac{1}{2}} P_2^{-\frac{1}{2}} P_1^{+\frac{1}{2}} \\
i{P_2^{-\frac{1}{2}}}_t &= {P_2^{-\frac{1}{2}}}_{xx} - 2P_2^{-\frac{1}{2}} Q^{+1} Q^{-1} - 2P_2^{+\frac{1}{2}} {Q^{-1}}_x + 2P_2^{+\frac{1}{2}} P_2^{-\frac{1}{2}} P_1^{-\frac{1}{2}}
\end{aligned} \tag{17}$$

3 AKNS Scheme with N=2 Superconformal Algebra (Neveu-Schwarz Type)

We generalize the connection given by Eq.(1) as

$$\Omega = \Omega_b + \Omega_f \tag{18}$$

where

$$\begin{aligned}
\Omega_b = \; & \Big(i\lambda J_0 + i\lambda L_0 + Q_1^{+m} J_{+m} + Q_1^{-m} J_{-m} + Q_2^{+m} L_{+m} + Q_2^{-m} L_{-m} \Big) dx + \\
& \Big(-A_1 J_0 - A_2 L_0 + B_1^{+m} J_{+m} + B_1^{-m} J_{-m} + B_2^{+m} L_{+m} + B_2^{-m} L_{-m} \Big) dt
\end{aligned} \tag{19}$$

$$\begin{aligned}
\Omega_f = \; & \Big(+P_1^{+\frac{m}{2}} G^1_{+\frac{m}{2}} + P_1^{-\frac{m}{2}} G^1_{-\frac{m}{2}} + P_2^{+\frac{m}{2}} G^2_{+\frac{m}{2}} + P_2^{-\frac{m}{2}} G^2_{-\frac{m}{2}} \Big) dx + \\
& \Big(+C_1^{+\frac{m}{2}} G^1_{+\frac{m}{2}} + C_1^{-\frac{m}{2}} G^1_{-\frac{m}{2}} + C_2^{+\frac{m}{2}} G^2_{+\frac{m}{2}} + C_2^{-\frac{m}{2}} G^2_{-\frac{m}{2}} \Big) dt
\end{aligned} \tag{20}$$

where L_0 , $J_{\pm m}$, $L_{\pm m}$ are bosonic generators and $G^1_{\pm\frac{m}{2}}$,$G^2_{\pm\frac{m}{2}}$ are fermionic generators of centerless N=2 superconformal algebra of Neveu-Schwarz type .In modern notation namely this algebra satisfy the following commutation and anticommutation relations

$$\begin{array}{lcl}
[L_r, L_s] & = & (r-s)\; L_{r+s} \\
[J_r, J_s] & = & 0 \\
[L_r, J_s] & = & -s\; J_{r+s} \\
\left\{G_r^1, G_s^2\right\} & = & 2\; L_{r+s} + (r-s)\; J_{r+s} \\
\left[J_r, G_s^{1,2}\right] & = & \pm G_{r+s}^{1,2} \\
\left[L_r, G_s^{1,2}\right] & = & (\frac{r}{2} - s)\; G_{r+s}^{1,2} \\
\left\{G_r^{1,2}, G_s^{1,2}\right\} & = & 0
\end{array} \tag{21}$$

Here, $J_{\pm m}$,$L_{\pm m}$ are generators with positive(negative) integer indices and $G^{1,2}_{\pm\frac{m}{2}}$ are generators with positive(negative) half integer indices. In Eq.(18) we assume summation over the repeated indices. The fields $Q_{1,2}^{\pm m}$ and $P_{1,2}^{\pm\frac{m}{2}}$ are x,t dependent and also functions $A_{1,2}$, $B_{1,2}^{\pm m}$ and $C_{1,2}^{\pm\frac{m}{2}}$ are x,t and λ dependent.

In N=2 superconformal algebra if we restrict $J_{\pm m}$ to have only J_0 components,$L_{\pm m}$ to have only L_0 , $L_{\pm 1}$ components, and $G^{1,2}_{\pm \frac{m}{2}}$ to have only $G^{1,2}_{\pm \frac{1}{2}}$ components we obtain sl(1,2) algebra given by Eq.(5) with the following definitions:

$$\begin{array}{l} J_1 = \frac{1}{2} J_0; \quad H_1 = -L_0; \quad E_{+1} = L_{+1}; \quad E_{-1} = -L_{-1} \\ F_{+\frac{1}{2}} = G^1_{+\frac{1}{2}}; \quad F_- = -\frac{1}{2} G^1_{-\frac{1}{2}}; \bar{F}_{+\frac{1}{2}} = \frac{1}{2} G^2_{+\frac{1}{2}}; \quad \bar{F}_{\frac{1}{2}} = -\frac{1}{2} G^2_{-\frac{1}{2}} \end{array} \tag{22}$$

From the integrability condition given by Eq.(6) we obtain

$$A_{1x} = \sum_{r=1}^{\infty} r(B_1^{-r} Q_2^{+r} - B_1^{+r} Q_2^{-r} + B_2^{-r} Q_1^{+r} - B_2^{+r} Q_1^{-r} - P_1^{-\frac{r}{2}} C_2^{+\frac{r}{2}} + P_1^{+\frac{r}{2}} C_2^{-\frac{r}{2}} + P_2^{-\frac{r}{2}} C_1^{+\frac{r}{2}} - P_2^{+\frac{r}{2}} C_1^{-\frac{r}{2}}) \tag{23}$$

$$A_{2x} = 2 \sum_{r=1}^{\infty} (r B_2^{-r} Q_2^{+r} - r B_2^{+r} Q_2^{-r} + P_1^{-\frac{r}{2}} C_2^{+\frac{r}{2}} + P_1^{+\frac{r}{2}} C_2^{-\frac{r}{2}} + P_2^{-\frac{r}{2}} C_1^{+\frac{r}{2}} + P_2^{+\frac{r}{2}} C_1^{-\frac{r}{2}}) \tag{24}$$

$$Q_1^{\pm m}{}_t = B_1^{\pm m}{}_x \mp i m \lambda B_1^{\pm m} \mp m A_2 Q_1^{\pm m} + \delta_J^{\pm m} \tag{25}$$

$$Q_2^{\pm m}{}_t = B_2^{\pm m}{}_x \mp i m \lambda B_2^{\pm m} \mp m A_2 Q_2^{\pm m} + \delta_L^{\pm m} \tag{26}$$

$$P_1^{\pm \frac{m}{2}}{}_t = C_1^{\pm \frac{m}{2}}{}_x \mp \tfrac{i}{2}(m \mp 2) \lambda C_1^{\pm \frac{m}{2}} \mp \tfrac{m}{2} P_1^{\pm \frac{m}{2}} A_2 + P_1^{\pm \frac{m}{2}} A_1 + \delta_{G^1}^{pmm} \tag{27}$$

$$P_2^{\pm \frac{m}{2}}{}_t = C_2^{\pm \frac{m}{2}}{}_x \mp \tfrac{i}{2}(m \pm 2) \lambda C_2^{\pm \frac{m}{2}} \mp \tfrac{m}{2} P_2^{\pm \frac{m}{2}} A_2 - P_2^{\pm \frac{m}{2}} A_1 + \delta_{G^2}^{\pm m} \tag{28}$$

where

$$\begin{aligned} \delta_J^{+m} = & \sum_{r,s=1}^{\infty} \Big[(r B_2^{+s} Q_1^{+r} - s B_1^{+s} Q_2^{+r}) \delta_{r+s,m} + (r-s)(P_1^{+\frac{r}{2}} C_2^{+\frac{s}{2}} - P_2^{+\frac{r}{2}} C_1^{+\frac{s}{2}}) \delta_{r+s,2m} \Big] \\ & + \sum_{\substack{r,s=1 \\ r>s}}^{\infty} \Big[(r B_2^{-s} Q_1^{+r} + s B_1^{-s} Q_2^{+r}) \delta_{r-s,m} + \frac{(r+s)}{2} (P_1^{+\frac{r}{2}} C_2^{-\frac{s}{2}} - P_2^{+\frac{r}{2}} C_1^{-\frac{s}{2}}) \delta_{r-s,2m} \Big] \\ & - \sum_{\substack{r,s=1 \\ r<s}}^{\infty} \Big[(r B_2^{+s} Q_1^{-r} + s B_1^{+s} Q_2^{-r}) \delta_{-r+s,m} + \frac{(r+s)}{2} (P_1^{-\frac{r}{2}} C_2^{+\frac{s}{2}} - P_2^{-\frac{r}{2}} C_1^{+\frac{s}{2}}) \delta_{-r+s,2m} \Big] \end{aligned} \tag{29}$$

$$\begin{aligned} \delta_L^{+m} = & \sum_{r,s=1}^{\infty} \Big[(r-s) B_2^{+s} Q_2^{+r} \delta_{r+s,m} + 2(P_1^{+\frac{r}{2}} C_2^{+\frac{s}{2}} + P_2^{+\frac{r}{2}} C_1^{+\frac{s}{2}}) \delta_{r+s,2m} \Big] \\ & + \sum_{\substack{r,s=1 \\ r>s}}^{\infty} \Big[(r+s) B_2^{-s} Q_2^{+r} \delta_{r-s,m} + 2(P_1^{+\frac{r}{2}} C_2^{-\frac{s}{2}} + P_2^{+\frac{r}{2}} C_1^{-\frac{s}{2}}) \delta_{r-s,2m} \Big] \\ & - \sum_{\substack{r,s=1 \\ r<s}}^{\infty} \Big[(r+s) B_2^{+s} Q_2^{-r} \delta_{-r+s,m} - 2(P_1^{-\frac{r}{2}} C_2^{+\frac{s}{2}} + P_2^{-\frac{r}{2}} C_1^{+\frac{s}{2}}) \delta_{-r+s,2m} \Big] \end{aligned} \tag{30}$$

$$\begin{aligned} \delta_{G^1}^{+m} = & -\frac{1}{2} \sum_{r,s=1}^{\infty} \Big[2B_1^{+s} - (r-s) B_2^{+s}) P_1^{+\frac{r}{2}} \delta_{r+2s,m} - 2(Q_1^{+r} + (r-s) Q_2^{+r}) C_1^{+\frac{s}{2}} \delta_{2r+s,m} \Big] \\ & + \frac{1}{2} \sum_{\substack{r,s=1 \\ 2r>s}}^{\infty} \Big[2Q_1^{+r} + (r+s) Q_2^{+r}) C_1^{-\frac{s}{2}} \delta_{2r-s,m} \Big] - \frac{1}{2} \sum_{\substack{r,s=1 \\ r>2s}}^{\infty} \Big[2(B_1^{-s} - (r+s) B_2^{-s}) P_1^{+\frac{r}{2}} \delta_{r-2s,m} \Big] \\ & - \frac{1}{2} \sum_{\substack{r,s=1 \\ r<2s}}^{\infty} \Big[2B_1^{+s} + (r+s) B_2^{+s}) P_1^{-\frac{r}{2}} \delta_{-r+2s,m} \Big] + \frac{1}{2} \sum_{\substack{r,s=1 \\ 2r<s}}^{\infty} \Big[2(Q_1^{-r} - (r+s) Q_2^{-r}) C_1^{+\frac{s}{2}} \delta_{-2r+s,m} \Big] \end{aligned} \tag{31}$$

$$\delta_{G^2}^{+m} = \delta_{G^1}^{+m} \left(\begin{array}{c} P_1 \rightarrow -P_2 \\ C_1 \rightarrow -C_2 \end{array} \right) \tag{32}$$

and

$$\delta_J^{-m}, \delta_L^{-m}, \delta_{G^1}^{-m}, \delta_{G^2}^{-m} = \delta_{G^1}^{+m}, \delta_{G^2}^{+m}, \delta_J^{+m}, \delta_L^{+m} \begin{pmatrix} +m \to -m \\ +r \to -r \\ +s \to -s \end{pmatrix} \tag{33}$$

In AKNS scheme we expand A_1,A_2,$B^{\pm 1}$,$C_1^{\pm\frac{1}{2}}$ and $C_2^{\pm\frac{1}{2}}$ in terms of the positive powers of λ as

$$A_1 = \sum_{n=0}^{2} \lambda^n a_{1n}; \quad A_2 = \sum_{n=0}^{2} \lambda^n a_{2n}; \quad B_1^{\pm m} = \sum_{n=0}^{2} \lambda^n b_{1n}^{\pm m}; \quad B_2^{\pm m} = \sum_{n=0}^{2} \lambda^n b_{2n}^{\pm m}; \tag{34}$$

$$C_1^{\pm\frac{m}{2}} = \sum_{n=0}^{2} \lambda^n c_{1n}^{\pm\frac{m}{2}}; \quad C_2^{\pm\frac{m}{2}} = \sum_{n=0}^{2} \lambda^n c_{2n}^{\pm\frac{m}{2}} \tag{35}$$

Inserting Eq.(34-35) into Eqs.(23-28)gives 30 relations in terms of a_{1n},a_{2n},$b_{1n}^{\pm m}$,$b_{2n}^{\pm m}$,$c_{1n}^{\pm\frac{m}{2}}$ and $c_{2n}^{\pm\frac{m}{2}}$ (n=0,1,2) . By solving these relations we get

$$a_{10} = i\sum_{r=1}^{\infty}(Q_2^{+r}Q_1^{-r} + Q_2^{-r}Q_1^{+r}) - i\sum_{r=1}^{\infty}\left[\frac{2}{r-2}\right]P_2^{-\frac{r}{2}}P_1^{+r} - i\sum_{r=1}^{\infty}\left[\frac{2}{r+2}\right]P_2^{+r}P_1^{-r}; \ a_{11} = a_{12} = 0;$$

$$a_{20} = 2i\sum_{r=1}^{\infty}Q_2^{+r}Q_2^{-r} - i\sum_{r=1}^{\infty}\left[\frac{4}{r-2}\right]P_2^{-r}P_1^{+r} + i\sum_{r=1}^{\infty}\left[\frac{4}{r+2}\right]P_2^{+r}P_1^{-r}; \ a_{21} = a_{22} = 0;$$

$$b_{10}^{\pm m} = \mp\frac{i}{m}Q_1^{\pm m}{}_x; \ b_{20}^{\pm m} = \mp\frac{i}{m}Q_2^{\pm m}{}_x; \ b_{11}^{\pm m} = Q_1^{\pm m}; \ b_{21}^{\pm m} = Q_2^{\pm m}; \ b_{12}^{\pm m} = 0; \ b_{22}^{\pm m} = 0 \tag{36}$$

$$c_{10}^{\pm\frac{m}{2}} = \mp\frac{2i}{m-2}P_1^{\pm\frac{m}{2}}{}_x; \ c_{20}^{\pm\frac{m}{2}} = \pm\frac{2i}{m+2}P_2^{\mp\frac{m}{2}}{}_x; \ c_{11}^{\pm\frac{m}{2}} = P_1^{\pm\frac{m}{2}}; \ c_{21}^{\pm\frac{m}{2}} = P_2^{\pm\frac{m}{2}}; \ c_{12}^{\pm\frac{m}{2}} = c_{22}^{\pm\frac{m}{2}} = 0$$

By using the relations given by Eq.(36) from Eqs.(25-28) we obtain the coupled super NLS equations as

$$\begin{aligned} -iQ_1^{\pm m}{}_t = &\mp\frac{1}{m}Q_1^{\pm m}{}_{xx} - 2Q_1^{\pm m}\left(\sum_{r=1}^{\infty}Q_2^{-r}Q_2^{+r}\right) \pm 4mQ_1^{\pm m}\left(\sum_{r=1}^{\infty}\left[\frac{1}{r-2}\right]P_2^{-\frac{r}{2}}P_1^{+\frac{r}{2}}\right) \\ &\mp 4mQ_1^{\pm m}\left(\sum_{r=1}^{\infty}\left[\frac{1}{r+2}\right]P_2^{+\frac{r}{2}}P_1^{-\frac{r}{2}}\right) + \delta_J^{\pm m} \end{aligned} \tag{37}$$

$$\begin{aligned} -iQ_2^{\pm m}{}_t = &\mp\frac{1}{m}Q_2^{\pm m}{}_{xx} - 2Q_2^{\pm m}\left(\sum_{r=1}^{\infty}Q_2^{-r}Q_2^{+r}\right) \pm 4mQ_2^{\pm m}\left(\sum_{r=1}^{\infty}\left[\frac{1}{r-2}\right]P_2^{-\frac{r}{2}}P_1^{+\frac{r}{2}}\right) \\ &\mp 4mQ_2^{\pm m}\left(\sum_{r=1}^{\infty}\left[\frac{1}{r+2}\right]P_2^{+\frac{r}{2}}P_1^{-\frac{r}{2}}\right) + \delta_L^{\pm m} \end{aligned} \tag{38}$$

$$\begin{aligned} -iP_1^{\pm\frac{m}{2}}{}_t = &\mp\frac{2}{m\mp 2}P_1^{\pm\frac{m}{2}}{}_{xx} \pm 2(m\mp 1)P_1^{\pm\frac{m}{2}}\left(\sum_{r=1}^{\infty}\left[\frac{1}{r-2}\right]P_2^{-\frac{r}{2}}P_1^{+\frac{r}{2}}\right) \\ &\mp 2(m\pm 1)P_1^{\pm\frac{m}{2}}\left(\sum_{r=1}^{\infty}\left[\frac{1}{r+2}\right]P_2^{+\frac{r}{2}}P_1^{-\frac{r}{2}}\right) + P_1^{\pm\frac{m}{2}}\sum_{r=1}^{\infty}Q_1^{+r}Q_2^{-r} \\ &+ P_1^{\pm\frac{m}{2}}\sum_{r=1}^{\infty}Q_1^{-r}Q_2^{+r} \mp mP_1^{\pm\frac{m}{2}}\sum_{r=1}^{\infty}Q_2^{+r}Q_2^{-r} + \delta_{G^1}^{\pm\frac{m}{2}} \end{aligned} \tag{39}$$

and

$$\begin{aligned}-P_2^{\pm\frac{m}{2}}{}_t = \mp\frac{2}{m\pm 2}P_2^{\pm\frac{m}{2}}{}_{xx} \pm 2(m\pm 1)P_2^{\pm\frac{m}{2}}\left(\sum_{r=1}^{\infty}\left[\frac{1}{r-2}\right]P_2^{-\frac{r}{2}}P_1^{+\frac{r}{2}}\right)\\ \mp 2(m\mp 1)P_2^{\pm\frac{m}{2}}\left(\sum_{r=1}^{\infty}\left[\frac{1}{r+2}\right]P_2^{+\frac{r}{2}}P_1^{-\frac{r}{2}}\right) - P_2^{\pm\frac{m}{2}}\sum_{r=1}^{\infty}Q_1^{+r}Q_2^{-r}\\ -P_2^{\pm\frac{m}{2}}\sum_{r=1}^{\infty}Q_1^{-r}Q_2^{+r} \mp mP_1^{\pm\frac{m}{2}}\sum_{r=1}^{\infty}Q_2^{+r}Q_2^{-r} + \delta_{G^2}^{\pm\frac{m}{2}}\end{aligned}\tag{40}$$

where

$$\begin{aligned}-i\delta_J^{+m} = \sum_{\substack{r=1\\r<m}}^{\infty}Q_2^{+r}Q_{1x}^{+(m-r)} - \sum_{\substack{r=1\\r<m}}^{\infty}\left[\frac{r}{m-r}\right]Q_1^{+r}Q_{2x}^{+(m-r)} + 2\sum_{\substack{r=1\\r<2m}}^{\infty}\left[\frac{r-m}{2m-r-2}\right]P_2^{+\frac{r}{2}}P_{1x}^{+(m-\frac{r}{2})}\\ -2\sum_{\substack{r=1\\r<2m}}^{\infty}\left[\frac{r-m}{2m-r+2}\right]P_1^{+\frac{r}{2}}P_{2x}^{+(m-\frac{r}{2})} + \sum_{\substack{r=1\\r>m}}^{\infty}Q_2^{+r}Q_{1x}^{-(r-m)} + \sum_{\substack{r=1\\r>m}}^{\infty}\left[\frac{r}{r-m}\right]Q_1^{+r}Q_{2x}^{-(r-m)}\\ +2\sum_{\substack{r=1\\r>2m}}^{\infty}\left[\frac{r-m}{r-2m-2}\right]P_1^{+\frac{r}{2}}P_{2x}^{-(\frac{r}{2}-m)} - 2\sum_{\substack{r=1\\r>2m}}^{\infty}\left[\frac{r-m}{r-2m+2}\right]P_2^{+\frac{r}{2}}P_{1x}^{-(\frac{r}{2}-m)} + \sum_{\substack{r=1\\r>m}}^{\infty}Q_2^{-(r-m)}Q_{1x}^{+r}\\ +\sum_{\substack{r=1\\r>m}}^{\infty}\left[\frac{(r-m)}{r}\right]Q_1^{-(r-m)}Q_{2x}^{+s} + 2\sum_{\substack{r=1\\r>2m}}^{\infty}\left[\frac{r-m}{r+2}\right]P_1^{-(\frac{r}{2}-m)}P_{2x}^{+\frac{s}{2}} - 2\sum_{\substack{r=1\\r>2m}}^{\infty}\left[\frac{r-m}{r-2}\right]P_2^{-(\frac{r}{2}-m)}P_{1x}^{+\frac{r}{2}}\end{aligned}\tag{41}$$

$$\begin{aligned}-i\delta_L^{+m} = -\sum_{\substack{r=1\\r<m}}^{\infty}\left[\frac{2r-m}{m-r}\right]Q_2^{+r}Q_{2x}^{+(m-r)} - 4\sum_{\substack{r=1\\r<2m}}^{\infty}\left[\frac{1}{2m-r+2}\right]P_1^{+\frac{r}{2}}P_{2x}^{+(m-\frac{r}{2})} - 4\sum_{\substack{r=1\\r<2m}}^{\infty}\left[\frac{1}{2m-r-2}\right]P_2^{+\frac{r}{2}}P_{1x}^{+(m-\frac{r}{2})}\\ +\sum_{\substack{r=1\\r>m}}^{\infty}\left[\frac{2r-m}{r-m}\right]Q_2^{+r}Q_{2x}^{-(r-m)} + 4\sum_{\substack{r=1\\r>2m}}^{\infty}\left[\frac{1}{r-2m+2}\right]P_1^{+\frac{r}{2}}P_{2x}^{+(\frac{r}{2}-m)} + 4\sum_{\substack{r=1\\r>2m}}^{\infty}\left[\frac{1}{r-2m+2}\right]P_2^{+\frac{r}{2}}P_{1x}^{-(\frac{r}{2}-m)}\\ +\sum_{\substack{r=1\\r>m}}^{\infty}\left[\frac{2r+m}{r}\right]Q_2^{-(r-m)}Q_{2x}^{+r} - 4\sum_{\substack{r=1\\r>2m}}^{\infty}\left[\frac{1}{r+2}\right]P_1^{-(\frac{r}{2}-m)}P_{2x}^{+\frac{r}{2}} - 4\sum_{\substack{r=1\\r>2m}}^{\infty}\left[\frac{1}{r-2}\right]P_2^{-(\frac{r}{2}-m)}P_{1x}^{+\frac{r}{2}}\end{aligned}\tag{42}$$

$$\begin{aligned}-i\delta_{G^1}^{+m} = \sum_{\substack{r=1\\2r<m}}^{\infty}\left[\frac{1}{r}\right]P_1^{+(\frac{m}{2}-r)}Q_{1x}^{+r} - \frac{1}{2}\sum_{\substack{r=1\\2r<m}}^{\infty}\left[\frac{m-3r}{r}\right]P_1^{+(\frac{m}{2}-r)}Q_{2x}^{+r} - 2\sum_{\substack{r=1\\2r<m}}^{\infty}\left[\frac{1}{m-2r-2}\right]P_{1x}^{+(\frac{m}{2}-r)}Q_1^{+r}\\ -2\sum_{\substack{r=1\\2r<m}}^{\infty}\left[\frac{3r-m}{m-2r-2}\right]P_{1x}^{+(\frac{m}{2}-r)}Q_2^{+r} + 2\sum_{\substack{r=1\\2r>m}}^{\infty}\left[\frac{1}{2r-m+2}\right]P_{1x}^{-(r-\frac{m}{2})}Q_1^{+r} + 2\sum_{\substack{r=1\\2r>m}}^{\infty}\left[\frac{3r-m}{2r-m+2}\right]P_{1x}^{-(r-\frac{m}{2})}Q_2^{+r}\\ -2\sum_{\substack{r=1\\r>m}}^{\infty}\left[\frac{1}{r-m}\right]P_1^{+\frac{r}{2}}Q_{1x}^{-(\frac{r}{2}-\frac{m}{2})} + \frac{1}{2}\sum_{\substack{r=1\\r>m}}^{\infty}\left[\frac{3r-m}{r-m}\right]P_1^{+\frac{r}{2}}Q_{2x}^{-(\frac{r}{2}-\frac{m}{2})} + \sum_{\substack{r=1\\2r>m}}^{\infty}\left[\frac{1}{r}\right]P_1^{-(r-\frac{m}{2})}Q_{1x}^{+r}\\ +\frac{1}{2}\sum_{\substack{r=1\\2r>m}}^{\infty}\left[\frac{3r-m}{r}\right]P_1^{-(r-\frac{m}{2})}Q_{2x}^{+r} - 2\sum_{\substack{r=1\\r>m}}^{\infty}\left[\frac{1}{r-2}\right]P_{1x}^{+\frac{r}{2}}Q_1^{-(\frac{r}{2}-\frac{m}{2})} - \frac{1}{2}\sum_{\substack{r=1\\r>m}}^{\infty}\left[\frac{3r-m}{r-2}\right]P_{1x}^{+\frac{r}{2}}Q_2^{-(\frac{r}{2}-\frac{m}{2})}\end{aligned}\tag{43}$$

$$\begin{aligned}-i\delta_{G^2}^{+m} = -\sum_{\substack{r=1\\2r<m}}^{\infty}\left[\frac{1}{r}\right]P_2^{+(\frac{m}{2}-r)}Q_{1x}^{+r} - \frac{1}{2}\sum_{\substack{r=1\\2r<m}}^{\infty}\left[\frac{m-3r}{r}\right]P_2^{+(\frac{m}{2}-r)}Q_{2x}^{+r} - 2\sum_{\substack{r=1\\2r<m}}^{\infty}\left[\frac{1}{m-2r+2}\right]P_{2x}^{+(\frac{m}{2}-r)}Q_1^{+r}\\ -2\sum_{\substack{r=1\\2r<m}}^{\infty}\left[\frac{3r-m}{m-2r+2}\right]P_{2x}^{+(\frac{m}{2}-r)}Q_2^{+r} + 2\sum_{\substack{r=1\\2r>m}}^{\infty}\left[\frac{1}{2r-m-2}\right]P_{2x}^{-(r-\frac{m}{2})}Q_1^{+r} + 2\sum_{\substack{r=1\\2r>m}}^{\infty}\left[\frac{3r-m}{2r-m-2}\right]P_{2x}^{-(r-\frac{m}{2})}Q_2^{+r}\end{aligned}$$

$$+2\sum_{\substack{r=1\\r>m}}^{\infty}\Big[\frac{1}{r-m}\Big]P_2^{+\frac{r}{2}}Q_{1x}^{-(\frac{r}{2}-\frac{m}{2})}+\frac{1}{2}\sum_{\substack{r=1\\r>m}}^{\infty}\Big[\frac{3r-m}{r-m}\Big]P_2^{+\frac{r}{2}}Q_{2x}^{-(\frac{r}{2}-\frac{m}{2})}-\sum_{\substack{r=1\\2r>m}}^{\infty}\Big[\frac{1}{r}\Big]P_2^{-(r-\frac{m}{2})}Q_{1x}^{+r} \tag{44}$$

$$+\frac{1}{2}\sum_{\substack{r=1\\2r>m}}^{\infty}\Big[\frac{3r-m}{r}\Big]P_2^{-(r-\frac{m}{2})}Q_{2x}^{+r}+2\sum_{\substack{r=1\\r>m}}^{\infty}\Big[\frac{1}{r+2}\Big]P_{2x}^{+\frac{r}{2}}Q_1^{-(\frac{r}{2}-\frac{m}{2})}+\frac{1}{2}\sum_{\substack{r=1\\r>m}}^{\infty}\Big[\frac{3r-m}{r-2}\Big]P_{2x}^{+\frac{r}{2}}Q_2^{-(\frac{r}{2}-\frac{m}{2})}$$

and

$$\delta_J^{-m},\delta_L^{-m},\delta_{G^1}^{-m},\delta_{G^2}^{-m}=\delta_{G^1}^{+m},\delta_{G^2}^{+m},\delta_J^{+m},\delta_L^{+m}\begin{pmatrix}+m\to -m\\ +r\to -r\end{pmatrix} \tag{45}$$

4 AKNS Scheme with N=2 Superconformal Algebra (Ramond Type)

We take the soliton connection as

$$\Omega=\Omega_b+\Omega_f \tag{46}$$

where

$$\begin{aligned}\Omega_b = &\Big(i\lambda L_0+Q_1^{+m}J_{+m}+Q_1^{-m}J_{-m}+Q_2^{+m}L_{+m}+Q_2^{-m}L_{-m}\Big)dx+\\ &\Big(-A_1J_0-A_2L_0+B_1^{+m}J_{+m}+B_1^{-m}J_{-m}+B_2^{+m}L_{+m}+B_2^{-m}L_{-m}\Big)dt\end{aligned} \tag{47}$$

$$\begin{aligned}\Omega_f = &\Big(+P_1^{+m}G^1_{+m}+P_1^{-m}G^1_{-m}+P_2^{+m}G^2_{+m}+P_2^{-m}G^2_{-m}\Big)dx+\\ &\Big(-A_3G^1_0-A_4G^2_0+C_1^{+m}G^1_{+m}+C_1^{-m}G^1_{-m}+C_2^{+m}G^2_{+m}+C_2^{-m}G^2_{-m}\Big)dt\end{aligned} \tag{48}$$

where L_0 , $J_{\pm m}$, $L_{\pm m}$ are bosonic generators and G_0^1,G_0^2, $G^1_{\pm m}$,$G^2_{\pm m}$ are fermionic generators of centerless N=2 superconformal algebra of Ramond type , namely they satisfy the following commutation and anticommutation relations

$$\begin{array}{lcl}[L_r,L_s] &=& (r-s)\ L_{r+s}\\ [J_r,J_s] &=& 0\\ [L_r,J_s] &=& -s\ J_{r+s}\\ \{G^1_r,G^2_s\} &=& 2\ L_{r+s}+(r-s)\ J_{r+s}\\ \big[J_r,G^{1,2}_s\big] &=& \pm G^{1,2}_{r+s}\\ \big[L_r,G^{1,2}_s\big] &=& (\frac{r}{2}-s)\ G^{1,2}_{r+s}\\ \big\{G^{1,2}_r,G^{1,2}_s\big\} &=& 0\end{array} \tag{49}$$

Here, $J_{\pm m}$,$L_{\pm m}$,$G^1_{\pm m}$ and $G^2_{\pm m}$ are generators with positive(negative) integer indices . In Eq.(46) we assume summation over the repeated indices. The fields $Q^{\pm m}_{1,2}$ and $P^{\pm m}_{1,2}$are x,t dependent and also functions $A_{1,2}$,$A_{3,4}$, $B^{\pm m}_{1,2}$ and $C^{\pm m}_{1,2}$ are x,t and λ dependent.

From the integrability condition given by Eq.(6) we obtain

$$A_{1x}=\sum_{r=1}^{\infty}r(B_1^{+r}Q_2^{-r}-B_1^{-r}Q_2^{+r}+B_2^{+r}Q_1^{-r}-B_2^{-r}Q_1^{+r}-2(P_1^{-r}C_2^{+r}-P_1^{+r}C_2^{-r}-P_2^{-r}C_1^{+r}+P_2^{+r}C_1^{-r})) \tag{50}$$

$$A_{2x}=\sum_{r=1}^{\infty}(2r(B_2^{-r}Q_2^{+r}-B_2^{+r}Q_2^{-r})+P_1^{-r}C_2^{+r}+P_1^{+r}C_2^{-r}+P_2^{-r}C_1^{+r}+P_2^{+r}C_1^{-r}) \tag{51}$$

$$A_{3x}=-\frac{3}{2}\sum_{r=1}^{\infty}r(B_2^{+r}P_1^{-r}-B_2^{-r}P_1^{+r}+C_1^{+r}Q_2^{-r}-C_1^{-r}Q_2^{+r})-(B_1^{+r}P_1^{-r}+B_1^{-r}P_1^{-r}-C_1^{+r}Q_1^{-r}-C_1^{-r}Q_1^{+r})) \tag{52}$$

$$A_{4x}=-\frac{3}{2}\sum_{r=1}^{\infty}r(B_2^{+r}P_2^{-r}-B_2^{-r}P_2^{+r}+C_2^{+r}Q_2^{-r}-C_2^{-r}Q_2^{+r})+(B_1^{+r}P_2^{-r}+B_1^{-r}P_2^{-r}-C_2^{+r}Q_1^{-r}-C_2^{-r}Q_1^{+r})) \tag{53}$$

$$Q_1^{\pm m}{}_t = B_1^{\pm m}{}_x \mp im\lambda B_1^{\pm m} \mp mP_1^{\pm m}A_4 \mp mP_2^{\pm m}A_3 \mp mA_2Q_1^{\pm m} + \delta_J^{\pm m} \tag{54}$$

$$Q_2^{\pm m}{}_t = B_2^{\pm m}{}_x \mp im\lambda B_2^{\pm m} - 2P_1^{\pm m}A_4 - 2P_2^{\pm m}A_3 \mp mA_2Q_2^{\pm m} + \delta_L^{\pm m} \tag{55}$$

$$P_1^{\pm m}{}_t = C_1^{\pm m}{}_x \mp im\lambda C_1^{\pm m} + P_1^{\pm m}A_1 \mp P_1^{\pm m}A_2 - A_3Q_1^{\pm m} \mp mA_3Q_2^{\pm m} + \delta_{G_1}^{\pm m} \tag{56}$$

$$P_2^{\pm m}{}_t = C_2^{\pm m}{}_x \mp im\lambda C_2^{\pm m} - P_1^{\pm m}A_1 \mp P_2^{\pm m}A_2 + A_4Q_1^{\pm m} \mp mA_4Q_2^{\pm m} + \delta_{G_2}^{\pm m} \tag{57}$$

where

$$\begin{aligned}
\delta_J^{+m} = &\sum_{r,s=1}^{\infty} \left((r-s)P_1^{+r}C_2^{+s} - (r-s)P_2^{+r}C_1^{+s} + rQ_1^{+r}B_2^{+s} - sQ_2^{+r}B_1^{+s}\right)\delta_{r+s,m} \\
&+ \sum_{\substack{r,s=1 \\ r>s}}^{\infty} \left((r+s)P_1^{+r}C_2^{-s} - (r+s)P_2^{+r}C_1^{-s} + rQ_1^{+r}B_2^{-s} + sQ_2^{+r}B_1^{-s}\right)\delta_{r-s,m} \\
&+ \sum_{\substack{r,s=1 \\ r<s}}^{\infty} \left((r+s)P_1^{-r}C_2^{+s} - (r+s)P_2^{-r}C_1^{+s} + rQ_1^{-r}B_2^{+s} + sQ_2^{-r}B_1^{+s}\right)\delta_{-r+s,m}
\end{aligned} \tag{58}$$

$$\begin{aligned}
\delta_L^{+m} = &\sum_{r,s=1}^{\infty} \left(2P_1^{+r}C_2^{+s} + 2P_2^{+r}C_1^{+s} + (r-s)Q_2^{+r}B_2^{+s}\right)\delta_{r+s,m} \\
&+ \sum_{\substack{r,s=1 \\ r>s}}^{\infty} \left(2P_1^{+r}C_2^{-s} + 2P_2^{+r}C_1^{-s} + (r+s)Q_2^{+r}B_2^{-s}\right)\delta_{r-s,m} \\
&+ \sum_{\substack{r,s=1 \\ r<s}}^{\infty} \left(2P_1^{-r}C_2^{+s} + 2P_2^{-r}C_1^{+s} - (r+s)Q_2^{-r}B_2^{+s}\right)\delta_{-r+s,m}
\end{aligned} \tag{59}$$

$$\begin{aligned}
\delta_{G^1}^{+m} = &- \sum_{r,s=1}^{\infty} \left(B_1^{+s}P_1^{+r} - \frac{1}{2}(2r-s)B_2^{+s}P_1^{+r} - C_1^{+s}Q_1^{+r} - \frac{1}{2}(r-2s)C_1^{+s}Q_2^{+r}\right)\delta_{r+s,m} \\
&- \sum_{\substack{r,s=1 \\ r>s}}^{\infty} \left(B_1^{-s}P_1^{+r} - \frac{1}{2}(2r+s)B_2^{-s}P_1^{+r} - C_1^{-s}Q_1^{+r} - \frac{1}{2}(r+2s)C_1^{-s}Q_2^{+r}\right)\delta_{r-s,m} \\
&- \sum_{\substack{r,s=1 \\ r<s}}^{\infty} \left(B_1^{+s}P_1^{-r} + \frac{1}{2}(2r+s)B_2^{+s}P_1^{-r} - C_1^{+s}Q_1^{-r} + \frac{1}{2}(r+2s)C_1^{+s}Q_2^{-r}\right)\delta_{-r+s,m}
\end{aligned} \tag{60}$$

$$\begin{aligned}
\delta_{G^2}^{+m} = &+ \sum_{r,s=1}^{\infty} \left(B_1^{+s}P_2^{+r} - \frac{1}{2}(2r-s)B_2^{+s}P_2^{+r} + C_2^{+s}Q_1^{+r} - \frac{1}{2}(r-2s)C_2^{+s}Q_2^{+r}\right)\delta_{r+s,m} \\
&+ \sum_{\substack{r,s=1 \\ r>s}}^{\infty} \left(B_1^{-s}P_2^{+r} - \frac{1}{2}(2r+s)B_2^{-s}P_2^{+r} + C_2^{-s}Q_1^{+r} - \frac{1}{2}(r+2s)C_2^{-s}Q_2^{+r}\right)\delta_{r-s,m} \\
&- \sum_{\substack{r,s=1 \\ r<s}}^{\infty} \left(B_1^{+s}P_2^{-r} + \frac{1}{2}(2r+s)B_2^{+s}P_2^{-r} + C_2^{+s}Q_1^{-r} + \frac{1}{2}(r+2s)C_2^{+s}Q_2^{-r}\right)\delta_{-r+s,m}
\end{aligned} \tag{61}$$

and

$$\delta_J^{-m}, \delta_L^{-m}, \delta_{G^1}^{-m}, \delta_{G^2}^{-m} = \delta_{G^1}^{+m}, \delta_{G^2}^{+m}, \delta_J^{+m}, \delta_L^{+m} \begin{pmatrix} +m \rightarrow -m \\ +r \rightarrow -r \\ +s \rightarrow -s \end{pmatrix} \tag{62}$$

In AKNS scheme we expand A_1,A_2,A_3,A_4,$B_1^{\pm m}$,$B_2^{\pm m}$,$C_1^{\pm m}$ and $C_2^{\pm m}$ in terms of the positive powers of λ as

$$A_1 = \sum_{n=0}^{2} \lambda^n a_{1n}; \quad A_2 = \sum_{n=0}^{2} \lambda^n a_{2n}; \quad A_3 = \sum_{n=0}^{2} \lambda^n a_{3n}; \quad A_4 = \sum_{n=0}^{2} \lambda^n a_{4n}; \tag{63}$$

$$B_1^{\pm m} = \sum_{n=0}^{2} \lambda^n b_{1n}^{\pm m};\quad B_2^{\pm m} = \sum_{n=0}^{2} \lambda^n b_{2n}^{\pm m};\quad C_1^{\pm m} = \sum_{n=0}^{2} \lambda^n c_{1n}^{\pm m};\quad C_2^{\pm m} = \sum_{n=0}^{2} \lambda^n c_{2n}^{\pm m} \tag{64}$$

Inserting Eq.(63-64) into Eqs.(50-57)gives 36 relations in terms of a_{1n},a_{2n},a_{3n},a_{4n},$b_{1n}^{\pm m}$,$b_{2n}^{\pm m}$, $c_{1n}^{\pm m}$ and $c_{2n}^{\pm m}$ (n=0,1,2) . By solving these relations we get

$$\begin{gathered}
a_{10} = i\sum_{r=1}^{\infty}(Q_1^{+r}Q_2^{-r} + Q_1^{-r}Q_2^{+r}) - 2i\sum_{r=1}^{\infty}(P_2^{-r}P_1^{+r} + P_2^{+r}P_1^{-r});\ a_{11} = a_{12} = 0; \\
a_{20} = 2i\sum_{r=1}^{\infty}(Q_2^{-r}Q_2^{+r}) - 2i\sum_{r=1}^{\infty}(\left[\frac{1}{r}\right]P_2^{-r}P_1^{+r} - \left[\frac{1}{r}\right]P_2^{+r}P_1^{-r});\ a_{21} = a_{22} = -i; \\
a_{30} = \frac{3}{2}i\sum_{r=1}^{\infty}(P_1^{+r}Q_2^{-r} + P_1^{-r}Q_2^{+r}) - i\sum_{r=1}^{\infty}\left[\frac{1}{r}\right]P_1^{+r}Q_1^{-r} + i\sum_{r=1}^{\infty}\left[\frac{1}{r}\right]P_1^{-r}Q_1^{+r};\ a_{31} = a_{32} = 0; \\
a_{40} = \frac{3}{2}i\sum_{r=1}^{\infty}(P_2^{+r}Q_2^{-r} + P_2^{-r}Q_2^{+r}) + i\sum_{r=1}^{\infty}\left[\frac{1}{r}\right]P_2^{+r}Q_1^{-r} - i\sum_{r=1}^{\infty}\left[\frac{1}{r}\right]P_2^{-r}Q_1^{+r};\ a_{41} = a_{42} = 0; \\
b_{10}^{\pm m} = \mp\frac{i}{m}{Q_1^{\pm m}}_x;\ b_{11}^{\pm m} = Q_1^{\pm m};\ b_{12}^{\pm m} = 0;\ b_{20}^{\pm m} = \mp\frac{i}{m}{Q_2^{\pm m}}_x;\ b_{21}^{\pm m} = Q_2^{\pm m};\ b_{22}^{\pm m} = 0 \\
c_{10}^{\pm m} = \mp\frac{i}{m}{P_1^{\pm m}}_x;\ c_{20}^{\pm m} = \pm\frac{i}{m}{P_2^{\mp m}}_x;\ c_{11}^{\pm m} = P_1^{\pm m};\ c_{21}^{\pm m} = P_2^{\pm m};\ c_{12}^{\pm m} = c_{22}^{\pm m} = 0
\end{gathered} \tag{65}$$

By using the relations given by Eq.(65) from Eqs.(54-57) we obtain the coupled super NLS equations as

$$\begin{gathered}
-i{Q_1^{\pm m}}_t = \pm\frac{1}{m}{Q_1^{\pm m}}_{xx} \pm \left(\sum_{r=1}^{\infty}P_2^{+r}Q_1^{-r}\right)P_1^{\pm m} \\
\pm P_2^{\pm m}\left(\sum_{r=1}^{\infty}P_1^{+r}Q_1^{-r}\right) \pm \left(\sum_{r=1}^{\infty}P_2^{-r}Q_1^{-r}\right)P_1^{\pm m} \mp P_2^{\pm m}\left(\sum_{r=1}^{\infty}P_1^{-r}Q_1^{+r}\right) \\
\mp 2\left(\sum_{r=1}^{\infty}P_2^{-r}P_1^{+r}\right)Q_1^{\pm m} \pm 2\left(\sum_{r=1}^{\infty}P_2^{+r}P_1^{-r}\right)Q_1^{\pm m} \pm 2m\left(\sum_{r=1}^{\infty}Q_2^{+r}Q_2^{-r}\right)Q_1^{\pm m} \\
\mp\frac{3}{2}m\left(\sum_{r=1}^{\infty}P_2^{+r}Q_2^{-r}\right)P_1^{\pm m} \pm \frac{3}{2}mP_2^{\pm m}\left(\sum_{r=1}^{\infty}P_1^{+r}Q_2^{-r}\right) \\
\mp\frac{3}{2}m\left(\sum_{r=1}^{\infty}P_2^{-r}Q_2^{+r}\right)P_1^{\pm m} \pm \frac{3}{2}mP_2^{\pm m}\left(\sum_{r=1}^{\infty}P_1^{-r}Q_2^{+r}\right) + \delta_J^{\pm m}
\end{gathered} \tag{66}$$

$$\begin{gathered}
-i{Q_2^{\pm m}}_t = \pm\frac{1}{m}{Q_2^{\pm m}}_{xx} - \frac{2}{m}\left(\sum_{r=1}^{\infty}P_2^{+r}Q_1^{-r}\right)P_1^{\pm m} \\
-\frac{2}{m}P_2^{\pm m}\left(\sum_{r=1}^{\infty}P_1^{+r}Q_1^{-r}\right) + \frac{2}{m}\left(\sum_{r=1}^{\infty}P_2^{-r}Q_1^{+r}\right)P_1^{\pm m} + \frac{2}{m}P_2^{\pm m}\left(\sum_{r=1}^{\infty}P_1^{-r}Q_1^{+r}\right) \\
-3\left(\sum_{r=1}^{\infty}P_2^{+r}Q_1^{-r}\right)P_1^{\pm m} + 3P_2^{\pm m}\left(\sum_{r=1}^{\infty}P_1^{+r}Q_2^{-r}\right) \\
-3\left(\sum_{r=1}^{\infty}P_2^{-r}Q_2^{+r}\right)P_1^{\pm m} + 3P_2^{\pm m}\left(\sum_{r=1}^{\infty}P_1^{-r}Q_2^{+r}\right) \mp 2\left(\sum_{r=1}^{\infty}P_2^{-r}P_1^{+r}\right)Q_2^{\pm m} \\
\pm 2\left(\sum_{r=1}^{\infty}P_2^{+r}P_1^{-r}\right)Q_2^{\pm m} \pm 2m\left(\sum_{r=1}^{\infty}Q_2^{+r}Q_2^{-r}\right)Q_2^{\pm m} + \delta_L^{\pm m}
\end{gathered} \tag{67}$$

$$
\begin{gathered}
-iP_1^{\pm m}{}_{t} = \pm\frac{1}{m}P_1^{\pm m}{}_{xx} + 4\left(\sum_{r=1}^{\infty} P_2^{+r}P_1^{+r}\right)P_1^{\pm m} \\
-\left(\sum_{r=1}^{\infty} Q_2^{-r}Q_1^{+r}\right)P_1^{\pm m} - \left(\sum_{r=1}^{\infty} Q_2^{+r}Q_1^{-r}\right)P_1^{\pm m} \pm 2m\left(\sum_{r=1}^{\infty} Q_2^{+r}Q_2^{-r}\right)P_1^{\pm m} \\
-\frac{1}{m}\left(\sum_{r=1}^{\infty} Q_1^{-r}P_1^{+r}\right)Q_1^{\pm m} + \frac{1}{m}\left(\sum_{r=1}^{\infty} Q_1^{+r}P_1^{-r}\right)Q_1^{\pm m} \\
+\frac{3}{2}\left(\sum_{r=1}^{\infty} Q_2^{-r}P_1^{+r}\right)Q_1^{\pm m} + \frac{3}{2}\left(\sum_{r=1}^{\infty} Q_2^{+r}P_1^{-r}\right)Q_1^{\pm m} \mp \frac{1}{2}\left(\sum_{r=1}^{\infty} Q_1^{-r}P_1^{+r}\right)Q_2^{\pm m} \\
\pm\frac{1}{2}\left(\sum_{r=1}^{\infty} Q_1^{+r}P_1^{-r}\right)Q_2^{\pm m} \pm \frac{3}{4}m\left(\sum_{r=1}^{\infty} Q_2^{-r}P_1^{+r}\right)Q_2^{\pm m} \pm \frac{3}{4}m\left(\sum_{r=1}^{\infty} Q_2^{+r}P_1^{-r}\right)Q_2^{\pm m} + \delta_{G^1}^{\pm m}
\end{gathered} \tag{68}
$$

$$
\begin{gathered}
-iP_2^{\pm m}{}_{t} = \pm\frac{1}{m}P_2^{\pm m}{}_{xx} - 4\left(\sum_{r=1}^{\infty} P_2^{-r}P_1^{+r}\right)P_2^{\pm m} \\
+\left(\sum_{r=1}^{\infty} Q_2^{-r}Q_1^{+r}\right)P_2^{\pm m} + \left(\sum_{r=1}^{\infty} Q_2^{+r}Q_1^{-r}\right)P_2^{\pm m} \pm 2m\left(\sum_{r=1}^{\infty} Q_2^{+r}Q_2^{-r}\right)P_2^{\pm m} \\
-\frac{1}{m}\left(\sum_{r=1}^{\infty} Q_1^{-r}P_2^{+r}\right)Q_1^{\pm m} + \frac{1}{m}\left(\sum_{r=1}^{\infty} Q_1^{+r}P_2^{-r}\right)Q_1^{\pm m} \\
-\frac{3}{2}\left(\sum_{r=1}^{\infty} Q_2^{-r}P_2^{+r}\right)Q_1^{\pm m} - \frac{3}{2}\left(\sum_{r=1}^{\infty} Q_2^{+r}P_2^{-r}\right)Q_1^{\pm m} \mp \frac{1}{2}\left(\sum_{r=1}^{\infty} Q_1^{-r}P_2^{+r}\right)Q_2^{\pm m} \\
\pm\frac{1}{2}\left(\sum_{r=1}^{\infty} Q_1^{+r}P_2^{-r}\right)Q_2^{\pm m} \pm \frac{3}{4}m\left(\sum_{r=1}^{\infty} Q_2^{-r}P_2^{+r}\right)Q_2^{\pm m} \pm \frac{3}{4}m\left(\sum_{r=1}^{\infty} Q_2^{+r}P_2^{-r}\right)Q_2^{\pm m} + \delta_{G^2}^{\pm m}
\end{gathered} \tag{69}
$$

where

$$
\begin{gathered}
-i\delta_J^{+m} = \sum_{r=1}^{\infty}\left[\frac{2r-m}{m-r}\right]P_1^{+r}P_{2x}^{+(m-r)} - \sum_{r=1}^{\infty}\left[\frac{r}{m-r}\right]P_2^{+r}P_{1x}^{+(m-r)} - \sum_{r=1}^{\infty} Q_2^{+r}Q_{1x}^{+(m-r)} \\
+\sum_{r=1}^{\infty}\left[\frac{r}{m-r}\right]Q_1^{+r}Q_{2x}^{+(m-r)} - \sum_{\substack{r=1\\ r>m}}^{\infty}\left[\frac{2r-m}{r-m}\right]P_1^{+r}P_{2x}^{-(r-m)} + \sum_{\substack{r=1\\ r>m}}^{\infty}\left[\frac{2r-m}{m-r}\right]P_2^{+r}P_{1x}^{-(r-m)} \\
-\sum_{\substack{r=1\\ r>m}}^{\infty} Q_2^{+r}Q_{1x}^{-(r-m)} + \sum_{\substack{r=1\\ r>m}}^{\infty}\left[\frac{r}{r-m}\right]Q_1^{+r}Q_{2x}^{-(r-m)} - \sum_{\substack{r=1\\ r<m}}^{\infty}\left[\frac{2r-m}{r}\right]P_1^{-(r-m)}P_{2x}^{+r} \\
+\sum_{\substack{r=1\\ r<m}}^{\infty}\left[\frac{2r-m}{r}\right]P_2^{-(r-m)}P_{1x}^{+r} - \sum_{\substack{r=1\\ r<m}}^{\infty} Q_2^{-(r-m)}Q_{1x}^{+r} - \sum_{\substack{r=1\\ r<m}}^{\infty}\left[\frac{r-m}{r}\right]Q_1^{+(r-m)}Q_{2x}^{+r}
\end{gathered} \tag{70}
$$

$$
\begin{gathered}
-i\delta_L^{+m} = \sum_{r=1}^{\infty}\left[\frac{2}{m-r}\right]P_1^{+r}P_{2x}^{+(m-r)} + \sum_{r=1}^{\infty}\left[\frac{2}{m-r}\right]P_2^{+r}P_{1x}^{+(m-r)} - \sum_{r=1}^{\infty}\left[\frac{2r-m}{m-r}\right]Q_2^{+r}Q_{1x}^{+(m-r)} \\
+\sum_{\substack{r=1\\ r>m}}^{\infty}\left[\frac{2}{r}\right]P_1^{+r}P_{2x}^{-(r-m)} + \sum_{\substack{r=1\\ r>m}}^{\infty}\left[\frac{2}{r-m}\right]P_2^{+r}P_{1x}^{-(r-m)} - \sum_{\substack{r=1\\ r>m}}^{\infty}\left[\frac{2r-m}{r-m}\right]Q_2^{+r}Q_{1x}^{-(r-m)} \\
+\sum_{\substack{r=1\\ r<m}}^{\infty}\left[\frac{2}{r}\right]P_1^{-(r-m)}P_{2x}^{+r} + \sum_{\substack{r=1\\ r<m}}^{\infty}\left[\frac{2}{r}\right]P_2^{-(r-m)}P_{1x}^{+r} - \sum_{\substack{r=1\\ r<m}}^{\infty}\left[\frac{2r-m}{r}\right]Q_2^{-(r-m)}Q_{1x}^{+r}
\end{gathered} \tag{71}
$$

$$
\begin{aligned}
-i\delta^{+m}_{G^1} = {} & \sum_{r=1}^{\infty}\Big[\frac{1}{m-r}\Big]Q_1^{+r}P_{1x}^{+(m-r)} + \sum_{r=1}^{\infty}\Big[\frac{1}{m-r}\Big]P_1^{+r}Q_{1x}^{+(m-r)} + \frac{1}{2}\sum_{r=1}^{\infty}\Big[\frac{3r-m}{m-r}\Big]Q_2^{+r}P_{1x}^{+(m-r)} \\
& +\frac{1}{2}\sum_{r=1}^{\infty}\Big[\frac{3r-m}{m-r}\Big]P_1^{+r}Q_{2x}^{+(m-r)} - \sum_{\substack{r=1\\ r>m}}^{\infty}\Big[\frac{1}{r-m}\Big]Q_1^{+r}P_{1x}^{-(r-m)} + \sum_{\substack{r=1\\ r>m}}^{\infty}\Big[\frac{1}{r-m}\Big]P_1^{+r}Q_{1x}^{-(r-m)} \\
& -\frac{1}{2}\sum_{\substack{r=1\\ r>m}}^{\infty}\Big[\frac{3r-2m}{r-m}\Big]Q_2^{+r}P_{1x}^{-(r-m)} - \frac{1}{2}\sum_{\substack{r=1\\ r>m}}^{\infty}\Big[\frac{3r-2m}{r-m}\Big]P_1^{+r}Q_{2x}^{-(r-m)} + \sum_{\substack{r=1\\ r<m}}^{\infty}\Big[\frac{1}{r}\Big]Q_1^{-(r-m)}P_{1x}^{+r} \\
& + \sum_{\substack{r=1\\ r<m}}^{\infty}\Big[\frac{1}{r}\Big]P_1^{-(r-m)}Q_{1x}^{+r} - \frac{1}{2}\sum_{\substack{r=1\\ r<m}}^{\infty}\Big[\frac{3r-m}{r}\Big]Q_2^{-(r-m)}P_{1x}^{+r} - \frac{1}{2}\sum_{\substack{r=1\\ r<m}}^{\infty}\Big[\frac{3r-m}{r}\Big]P_1^{-(r-m)}Q_{2x}^{+r}
\end{aligned} \tag{72}
$$

$$
\begin{aligned}
-i\delta^{+m}_{G^2} = {} & -\sum_{r=1}^{\infty}\Big[\frac{1}{m-r}\Big]Q_1^{+r}P_{2x}^{+(m-r)} + \sum_{r=1}^{\infty}\Big[\frac{1}{m-r}\Big]P_1^{+r}Q_{1x}^{+(m-r)} + \frac{1}{2}\sum_{r=1}^{\infty}\Big[\frac{3r-m}{m-r}\Big]Q_2^{+r}P_{2x}^{+(m-r)} \\
& +\frac{1}{2}\sum_{r=1}^{\infty}\Big[\frac{3r-m}{m-r}\Big]P_1^{+r}Q_{2x}^{+(m-r)} + \sum_{\substack{r=1\\ r>m}}^{\infty}\Big[\frac{1}{r-m}\Big]Q_1^{+r}P_{2x}^{-(r-m)} + \sum_{\substack{r=1\\ r>m}}^{\infty}\Big[\frac{1}{r-m}\Big]P_1^{+r}Q_{1x}^{-(r-m)} \\
& -\frac{1}{2}\sum_{\substack{r=1\\ r>m}}^{\infty}\Big[\frac{3r-2m}{r-m}\Big]Q_2^{+r}P_{2x}^{-(r-m)} - \frac{1}{2}\sum_{\substack{r=1\\ r>m}}^{\infty}\Big[\frac{3r-2m}{r-m}\Big]P_1^{+r}Q_{2x}^{-(r-m)} - \sum_{\substack{r=1\\ r<m}}^{\infty}\Big[\frac{1}{r}\Big]Q_1^{-(r-m)}P_{2x}^{+r} \\
& + \sum_{\substack{r=1\\ r<m}}^{\infty}\Big[\frac{1}{r}\Big]P_1^{-(r-m)}Q_{1x}^{+r} - \frac{1}{2}\sum_{\substack{r=1\\ r<m}}^{\infty}\Big[\frac{3r-m}{r}\Big]Q_2^{-(r-m)}P_{2x}^{+r} - \frac{1}{2}\sum_{\substack{r=1\\ r<m}}^{\infty}\Big[\frac{3r-m}{r}\Big]P_1^{-(r-m)}Q_{2x}^{+r}
\end{aligned} \tag{73}
$$

and

$$
\delta_J^{-m}, \delta_L^{-m}, \delta_{G^1}^{-m}, \delta_{G^2}^{-m} = \delta_{G^1}^{+m}, \delta_{G^2}^{+m}, \delta_J^{+m}, \delta_L^{+m} \left(\begin{array}{c} +m \rightarrow -m \\ +r \rightarrow -r \end{array} \right) \tag{74}
$$

5 Conclusions

Using AKNS scheme and N=2 superconformal algebra of Neveu-Schwarz and Ramond types we obtain two different new super- extensions of coupled Nonlinear Schrödinger equations.These super- extensions of coupled NLS equations involve infinite number of bosonic and infinite number of fermionic fields,and mathematically my work is an extension of the work done in ref.6. The solution of these NLS equations with infinite number of bosonic and infinite number of femionic fields is an open problem.

References

[1] Ablowitz MJ,Kaub JD, Newell AC,Segur H. Phys Rev Lett 1973;30:1262;
Ablowitz MJ,Kaub JD, Newell AC,Segur H. Phys Rev Lett 1973;31:125;
Ablowitz MJ,Kaub JD, Newell AC,Segur H. Stud App Math 1974;53:249.

[2] Fordy AP,Kulish PP. Commun Math Phys 1983;89:427.

[3] Gürses M, Oğuz Ö,Salihoğlu S. Int Mod Phys A 1990;5:1801.

[4] Canoğlu A, Güldoğan B, Salihoğlu S. Int Mod Phys A 1991;6:4655;
Canoğlu A, Güldoğan B, Salihoğlu S. Int Mod Phys A 1992;7:7287;

[5] Özer HT, Salihoğlu S. Chaos, Solitons and Fractals 2006;30:363.

[6] Özer HT, Salihoğlu S. Chaos, Solitons and Fractals 2007;33:1417.

[7] Ademollo M,Brink L ,d'Adda A,d'Auria R, Napolitano E,Sciuto S, et al. Phys Lett B 1976;62:105.

[8] Kac VG. Adv Math 1977;26:8.

[9] Ademollo M,at al. Nucl Phys B 1976;111:77;
Ademollo M,at al. Nucl Phys B 1976;114:297.

[10] Brink L, Schwarz JH. Nucl Phys B 1977;121:285

[11] Fradkin ES, Tseytlin AA. Phys Lett B 1981;106:63;
Fradkin ES, Tseytlin AA. Phys Lett B 1985;162:295.

[12] Mathur SD, Mukhi S. Phys Rev D 1987;36:465;
Mathur SD, Mukhi S. Nucl Phys B 1988;302:130.

[13] Ouguri H, Vafa C. Nucl Phys B 1991;361:469;
Ouguri H, Vafa C. Nucl Phys B 1991;367:83.

[14] Martinec E.hep-th/9710122.

[15] Gato-Rivera B, Semikhatov AM. Phys Lett B 1992;293:72;
Gato-Rivera B, Semikhatov AM. Nucl Phys B 1993;408:133.

[16] Bershadsky M, Lerche W, Nemeschansky D, Warner NP. Nucl Phys B 1993;401:304.

[17] Roelofs GHM, Kersten PHM. J. Math. Phys. 1992;33:2185.

[18] Brunelli JC,Das A. J.Math.Phys. 1995;36:268.

[19] Popowicz Z. Phys. Lett A 1994;194:375.